
Buckeye Valley Local School District Self Medication for Rescue Inhaler Authorization Form

An Asthma Action Plan MUST accompany this form


		Date of Birth					
Student Address:							
School:			ox/city/zip code)		grade:		
Reason for carrying R If other, please specify				Peanuts	Tree Nuts	Other	
	edication: Dose:				frequency:		
Start date:		End D	ate:				
Adverse reactions that	should be repo	orted to the	e physician :				
Severe Adverse reacti					not prescribed, s	should they	
Other instructions:							
By signing below, the above student carry the the student is self-adm preventing adults from use of the rescue inhallother student.	e rescue inhale ninistering medi responding ap	er on their cation, no propriately	person at school a adult may be awar in an emergency;	nd any school fur e that the studen and that the stud	nction. They rea t is experiencino lent has been fu	alize that because g difficulty, ully trained in the	
In the event that the re responsibility to assum for the student, classm	ne control of the	rescue in	-		-		
Physician Name:				Phone numb	oer:		
Physician Signature:_				Date:			
Parent/Guardian Name	e:			Phone Num	ber:		
Parent/Guardian Signa	ature:			Date:			
By signing below, the condition, is able to ide of technique of admini agrees to NEVER sha IMMEDIATELY in the	entify signs/ syr stering medicat re the rescue ir	nptoms of ion using ihaler with	respiratory distres a rescue inhaler w n another person. T	s and identify the ith another perso he student agree	need for inhale n. The above n es to seek adult	r use and mastery amed student help	
Student Signature:				Date:		·	

Buckeye Valley Local School District

How to Use a Metered-Dose Inhaler with an Aerochamber (Spacer)




1. Shake the medicine.




Insert the mouthpiece of the inhaler into the rubber-sealed end of the Aerochamber.



 Breathe all the air out of your lungs then put the Aerochamber into your mouth between your teeth. Make a tight seal around the mouthpiece with your lips.



 Press the metered-dose inhaler down once to release a spray of medicine.
 The medicine will be trapped in the spacer. Breathe in slowly and deeply.



Hold your breath for at least 5 to 10 seconds. Breathe out slowly.

Copyright @2001 McKesson Health Solutions LLC. All rights reserved.